

## poladay CP

35% carbamide peroxide

## advanced tooth whitening system INSTRUCTIONS FOR USE FOR PATIENTS AND DENTAL PROFESSIONALS

Pola Day CP is a 35% carbamide peroxide, high viscosity, neutral pH take home tooth whitening gel. It is a pleasantly flavoured, crystal clear sustained release gel with a water content high enough to reduce sensitivity.

INDICATIONS FOR USE:

Whitening of discoloured vital teeth.

Whitening of discoloured non-vital teeth.

COMPOSITION: 35% Carbamide peroxide 40% Water 0.5% Flavour 24.5% Thickening agents

**NET WEIGHT:**Depending on kit purchased, 1.3g syringes or 3g syringes Please refer to outside of box.

IYDROGEN PEROXIDE EQUIVALENT: ola Day CP 35% Carbamide Peroxide = 11.7% Hydrogen eroxide

- INSTRUCTIONS:
   Place a small drop of gel into every compartment of the tray for the teeth undergoing treatment.
   Seat the tray, with the gel around the teeth.
   Wipe away excess gel in mouth with a tissue or dry soft brush.
   After 15 minutes of treatment, remove tray. Rinse tray and mouth with lukewarm water.
   Note: Treatment time can be up to 30 minutes if required.
   Brush teeth.





3. 5





# -OFFICE JUMP START (FOR DENTAL PROFESSIONALS) fren used as a jump start in the dental office, treatment tim 30 minutes.

- TORAGE AND HANDLING: Store product in the refrigerator (2°-8°C / 35°-45°F). Do not use after expiration date. Use product at room temperature.
- RECAUTIONS:

- To be used under the supervision of a dentist.
  Do not eat, drink or smoke during treatment.
  Do not smoke immediately after treatment wait for at

- Do not smoke immediately after treatment wait for at least two hours.

  Food and drinks containing strong colours should be avoided or consumed in moderation during the entire treatment period.

  Keep out of reach of children and pets.

  Not to be used by pregnant or lactating women or children under 14 years old.

  Patients with a history of chemical allergies are advised to carry out allergy testing by a specialist before using this norduct. Patients with a history of chemical allergies are advised to carry out allergy testing by a specialist before using this product.
  Discontinue use if any unusual sensitivity or reactions occur and consult your dentist.
  Do not inject Pola Day CP into the body.
  CAUTION: FEDERAL LAW RESTRICTS THIS DEVICE TO SALE BY OR ON THE ORDER OF A DENTIST.
  MSDS available at www.sdi.com.au or contact your regional representative.

- RST AID: Eye (contact): Wash thoroughly with water and seek medical advice if symptoms persist. Skin/Tissue (contact): Wash thoroughly with water and seek medical advice if symptoms persist. Ingestion (large amount): Drink plenty of milk or water and seek medical advice.







Last Revision : 2012 - 08