## CREATED BY SDI



## STEP BY STEP GUIDE: TAKE HOME

Firstly, brush teeth as normal. Now, remove the whitening gel syringe cap and insert an application tip by twisting it securely onto the syringe.

- Place a small drop of gel into every compartment of the tray for the teeth undergoing treatment.
  Tip: Use the markings on the syringe as a guide:
  3g syringe - For each whitening application, extrude approximately one sixth (0.5g) of the syringe on each of the top and bottom whitening trays.
  1.3g syringe - For each whitening application, extrude approximately one third (0.4g) of the syringe on each of the top and bottom whitening trays.
- 2. Seat the tray with the gel, around your teeth.
- **3.** Wipe away any excess or overflowing gel with a tissue or dry soft brush.
- **4.** After treatment, remove your tray. Rinse tray and mouth with lukewarm water.
- 5. Brush teeth.





## PLEASE NOTE:

- Use gel at room temperature
- Do not eat, drink or smoke during treatment
- Do not smoke immediately after treatment wait for at least two hours
- Foods and drinks containing strong colours should be avoided for at least 48 hours or consumed in moderation
- Not to be used by pregnant or lactating women
- · Certain countries have age restrictions for whitening products where it is not to be used under a certain age

## TREATMENT GUIDELINES

	6% & 7.5%		9.5%
POLA DAY	l or 2 x 30 minutes / day or l x 45minutes / day		2 x 15 minutes/day or 1 x 30 minutes / day
POLA NIGHT	10%	16% & 18%	22%
	1 x 2 hours / day to overnight	1 x 90 minutes / day to overnight	1 x 45 minutes / day

ADVANCED TOOTH WHITENING SYSTEM

> YOUR **our** Smile **Vision**.







